

The Kinetic Flux of Quasar Jets

Daniel A. Schwartz¹

**Herman L. Marshall², Jonathan Gelbord², Diana M. Worrall^{1,3}, Mark Birkinshaw^{1,3},
James E. J. Lovell⁴, David L. Jauncey⁴, Eric S. Perlman⁵, David W. Murphy⁶**

1. Harvard/Smithsonian Center for Astrophysics

2. Massachusetts Institute of Technology

3. University of Bristol

4. Australia Telescope National Facility, CSIRO

5. University of Maryland Baltimore Campus

6. Jet Propulsion Laboratory



Summary

X-Ray Observations Provide Evidence that Radio Jets:

- 1. Carry Large Flux of Energy – Greater than Accretion Luminosity**
- 2. Transport Energy Very Efficiently – $L_{jet} / K_{jet} \lesssim 10^{-5}$**

We Predict:

- 3. X-ray Jets become more prominent at large redshift**

Outline


1. X-Ray Observations of Radio Jets

2. Spatially Resolved Analysis

3. Spectral Energy Distribution

4. Interpretation as IC/CMB

5. The Kinetic Power



Outline






1. X-Ray Observations

2. Spatially Resolved Analysis

3. Spectral Energy Distribution

4. Interpretation as IC/CMB

5. The Kinetic Power



Outline





1. X-Ray Observations

2. Spatially Resolved Analysis

3. Spectral Energy Distribution

4. Interpretation as IC/CMB

5. The Kinetic Power



Outline





1. X-Ray Observations

2. Spatially Resolved Analysis

3. Spectral Energy Distribution

4. Interpretation as IC/CMB

5. The Kinetic Power



Outline


1. X-Ray Observations

2. Spatially Resolved Analysis


3. Spectral Energy Distribution

4. Interpretation as IC/CMB


5. The Kinetic Power




PKS 0920-397 K 1





PKS 0920-397 K 1



PKS 0920-397 K 1







The intersection gives a solution for the magnetic field, B , in the rest frame, and for the apparent Doppler factor,

$$\delta = (\Gamma(1 - \beta \cos(\theta)))^{-1}.$$


Here, $B=9.6 \mu\text{G}$ and $\delta=6.6$.

Uncertainties in the Magnetic Field Estimates

Uncertainties in the Magnetic Field Estimates


Uncertainties in the Magnetic Field Estimates

Equipartition





Uncertainties in the Magnetic Field Estimates

Equipartition




Inverse Compton





- **Determined B and δ within a factor of 2**
- **Can reduce IC region, depending on precision of radio spectral index**
- **Kinetic flux is $\propto (B\delta)^2$, for equipartition**


Structure of the Jets



Kinetic Flux

From $K = \Gamma^2 \pi r^2 \beta c U,$

$K \propto \delta^2 \theta_r^2 (3 B^2 / (8 \pi))$



PKS 0208-512

PKS 0920-397


PKS 1030-357

PKS 1202-512

Kinetic Flux

From $K = \Gamma^2 \pi r^2 \beta c U,$

$$K \propto \delta^2 \theta_r^2 (3 B^2 / (8 \pi))$$



PKS 0208-512

PKS 0920-397


PKS 1030-357


PKS 1202-512

Kinetic Flux

Energy in Protons?

- Large symbols assume $U_p = U_e$
- Lower symbols assume pure e^\pm plasma
- Upper symbols assume cold protons, $n_p = n_e$, and $\langle \gamma \rangle_e = 183$







Implications of the X-Ray Jets

- **Eddington Luminosity might not limit Accretion Rate**
- **Jets may Power Cluster Cavities – Stop Cooling Flows**
- **IC/CMB X-ray jets Maintain Constant Surface Brightness vs. z . We will detect them at Arbitrarily Large Redshift.**

Where ARE the bright X-ray Jets at High Redshift?



Gurvits et al. 2003

Where ARE the bright X-ray Jets at High Redshift?

HERE!




Siemiginowska et al. 2003

There Should Be Radio Quiet X-Ray Jets!

- **1 keV X-rays produced by $\gamma \approx 1000/\Gamma$**
- **$\nu = 4.2 \times 10^{-6} \gamma^2 \text{ H}[\mu\text{G}] \approx 10 \text{ MHz}$**


A Radio Quiet X-Ray Jet?

EMSS 0841+1314



← 30" →


J0841





30" = 273 kpc
← →

A Radio Quiet X-Ray Jet?


EMSS 0841+1314



30"




J0841




30" = 273 kpc

A Radio Quiet X-Ray Jet?


EMSS 0841+1314





30"



J0841



30" = 273 kpc



Summary

X-Ray Observations Provide Evidence that Radio Jets:

- 1. Carry Large Flux of Energy – Greater than Accretion Luminosity**
- 2. Transport Energy Very Efficiently – $L_{jet} / K_{jet} \lesssim 10^{-5}$**

We Predict:

- 3. X-ray Jets become more prominent at large redshift**